

Literature Review and Analysis: Coastal Urbanization and Microbial Contamination of Shellfish Growing Areas

Stuart Glasoe, Puget Sound Action Team and Aimee Christy, Pacific Shellfish Institute


Introduction

Many factors influence the suitability of coastal areas for growing and harvesting shellfish, and none is more vital than clean water. Human habitation has had a dramatic effect on the condition of the nation's coastal habitats and resources. A primary concern in shellfish growing areas is microbial contamination from human and animal feces. Key sources of fecal pollution include discharges from municipal sewage treatment plants, on-site sewage systems, stormwater runoff, marinas and boaters, farm animals, pets and wildlife.

The condition and classification of shellfish growing areas tend to correlate with population densities and development levels in adjacent watersheds, but our limited understanding of these relationships hampers efforts to permanently safeguard water quality for shellfish harvesting. To better understand and address these issues, the Puget Sound Action Team initiated a study to assess the effects of urbanization on shellfish growing areas that included a literature review and analysis, completed in June 2004.



Elliott Bay, King County — closed to harvest.



Totten Inlet, Thurston County — open to harvest.

Figure 1. Are there measurable correlations between coastal development and nearshore bacterial levels? If so, what are the mechanisms that drive these relationships? Which landscape indicators correlate most strongly with nearshore bacterial levels?

Effects of Urbanization on Watershed Hydrology and Water Quality

Numerous studies have documented the effects of development on the condition of streams, nearshore habitats and other aquatic ecosystems. Impacts include the fragmentation and loss of habitat and the degradation of water resources and water quality.

“Of all the land use changes affecting the hydrology of an area, urbanization is by far the most forceful” (Leopold 1968). A central issue is the change in flow regime, particularly the loss of water storage in the soil column in the Pacific Northwest's native rain forests (Booth 2000). Effects begin with the first expression of human activity in a watershed and then progress as development increases in scope and scale (Booth *et al.* 2001). Clearing of land cover and construction of impervious surfaces converts large volumes of water from subsurface flows to surface runoff. The landscape's capacity to attenuate flows and break down pollutants is further reduced as other features of the terrain are ditched, drained, piped and armored to efficiently shed runoff. This combination of reduced retention and enhanced conveyance causes increased flooding, degraded stream channels, reduced ground-water recharge, lower stream baseflows, and polluted surface waters and shellfish beds.




Figure 2. Population densities and commercial shellfish growing area classifications in Puget Sound. Shellfish areas not officially classified are closed to commercial harvest, including the Everett-to-Tacoma urban corridor.

Selected Studies of Urbanization and Shellfish Contamination

Mailo and Tschetter 1981. Correlated population increases in two North Carolina counties with degraded water quality and shellfish closures. Attributed impacts mainly to growth that had outstripped the region's sewage management capacity.

Duda and Cromartie 1982. Documented sharp increases in development and shellfish closures in coastal North Carolina. Correlated bacterial levels with densities of on-site sewage systems and identified stormwater runoff and drainage modifications as contributing factors.

Macfarlane 1996. Attributed shellfish closures in Cape Cod, Massachusetts, with rapid development and loadings from stormwater, on-site sewage systems and wildlife. Restoration efforts focused on installation of stormwater treatment systems.

Weiskel *et al.* 1996. Assessed bacterial sources, pathways and loadings in Massachusetts' Buttermilk Bay watershed. Determined that stormwater runoff contributed nearly a quarter of the bacterial load, had a disproportionately high impact on nearshore water quality, and correlated strongly with urban land uses.

White *et al.* 2000. Correlated bacterial loads and shellfish closures in the lightly developed Jumping Run Creek watershed of North Carolina with ditching, bulkheading and other hydrologic modifications that enhanced runoff and drainage.

Mallin *et al.* 2000, 2001. Correlated bacterial levels and shellfish closures in North Carolina tidal creeks with population growth on a regional scale, and with population, developed land, and impervious cover on a watershed scale. Further analysis highlighted the importance of limited impervious cover and intact land cover and wetland systems in mitigating coastal microbial contamination.

Lipp *et al.* 2001a, 2001b. Correlated concentrations and spatial and seasonal distribution of bacterial loadings with freshwater

flows and densities of on-site sewage systems in Florida's Charlotte Harbor and Sarasota Bay. Identified subsurface flows, surface runoff and tides as key physical factors.

Kelsey *et al.* 2003, 2004. Employed a variety of techniques to compare land uses and ecosystem responses in urban Murrells Inlet and rural North Inlet of South Carolina. Attributed greater bacterial contamination and differences in bacterial profiles in Murrells Inlet to urban influences, primarily stormwater runoff.




Figure 3. Sample results correlating impervious cover and bacterial levels in North Carolina tidal creeks (Mallin *et al.* 2001).

Conclusions

■ Two significant and related trends—population growth and urbanization—are stressing and degrading coastal ecosystems.

■ Bivalve molluscan shellfish are efficient vectors of enteric viruses and other pathogens. Actions that prevent contamination of shellfish growing areas are vital for safeguarding public health.

■ Microbial contamination is chronic in many coastal areas of the country and correlates with population densities, development levels, rainfall events, stormwater runoff and river flows.

■ Urbanization dramatically reduces the capacity of watersheds to attenuate flows and contaminants. The imprint of urbanization is generally permanent and the related impacts are equally difficult to mitigate or reverse.

■ Pollution impacts are exacerbated by development practices that disrupt hydrologic processes and increase connectivity between pollution sources and downstream waters.

■ Research correlating development with degradation of stream systems is extensive and compelling. Research on nearshore



Figure 4. Piped stormwater discharging to Puget Sound shellfish waters.

marine systems, including shellfish growing areas, is more limited but reveals strong and similar relationships.

■ Impervious cover is the most widely researched landscape indicator for gauging the effects of development. Both freshwater and marine systems exhibit a continuum of effects that surface at low levels of development (<10 percent impervious cover) and then amplify as development progresses.

■ The scientific literature supports long-standing observations that urban development is incompatible with safe shellfish harvesting, but does not reveal a universal rule or threshold for determining suitable land uses and development levels.

■ Pollution impacts can be mitigated, but not eliminated, using a variety of approaches. Sound land use planning, pollution prevention, watershed protection and personal stewardship are needed to guide appropriate development practices.

Literature Cited

- Booth, D.B. 2000. *Forest Cover, Impervious-Surface Area, and the Mitigation of Urbanization Impacts in King County, Washington*. Prepared for King County Water and Land Resources Division. Seattle, Washington. 18 pp.
- Booth, D.B., J.R. Karr, S. Schauman, C.P. Konrad, S.A. Morley, M.G. Larson, P.C. Henshaw, E.J. Nelson and S.J. Burges. 2001. *Urban Stream Rehabilitation in the Pacific Northwest: Physical, Biological and Social Considerations*. Final Report, EPA Grant Number R82-5284-010. University of Washington. Seattle, Washington. 78 pp.
- Duda, A.M. and K.D. Cromartie. 1982. Coastal Pollution from Septic Tank Drainfields. *Journal of the Environmental Engineering Division, American Society of Civil Engineers*. 108(6):1265-1279.
- Kelsey, H., G. Scott, D.E. Porter, B. Thompson and L. Webster. 2003. Using Multiple Antibiotic Resistance and Land Use Characteristics to Determine Sources of Fecal Coliform Bacterial Pollution. *Environmental Monitoring and Assessment*. 81(1-3):337-348.
- Kelsey, H., D.E. Porter, G. Scott, M. Neet and D. White. 2004. Using Geographic Information Systems and Regression Analysis to Evaluate Relationships Between Land Use and Fecal Coliform Bacterial Pollution. *Journal of Experimental Marine Biology and Ecology*. 298(2):197-209.
- Leopold, L.B. 1968. *Hydrology for Urban Land Planning—A Guidebook on the Hydrologic Effects of Urban Land Use*. U.S. Geological Survey Circular 554. Washington, D.C. 18 pp.
- Lipp, E.K., S.A. Farrah and J.B. Rose. 2001a. Assessment and Impact of Microbial Fecal Pollution and Human Enteric Pathogens in a Coastal Community. *Marine Pollution Bulletin*. 42(4):286-293.
- Lipp, E.K., R. Kurz, R. Vincent, C. Rodriguez-Palacios, S.R. Farrah and J.B. Rose. 2001b. The Effects of Seasonal Variability and Weather on Fecal Pollution and Enteric Pathogens in a Subtropical Estuary. *Estuaries*. 24(2): 266-276.
- Macfarlane, S.L. 1996. Shellfish as the Impetus for Embayment Management. *Estuaries*. 19(2A): 311-319.
- Maiolo, J. and P. Tschetter. 1981. Relating Population Growth to Shellfish Bed Closures: A Case Study from North Carolina. *Coastal Zone Management Journal*. 9(1):1-18.
- Mallin, M.A., S.H. Ensign, M.R. McIver, G.C. Shank and P.K. Fowler. 2001. Demographic, Landscape, and Meteorological Factors Controlling the Microbial Pollution of Coastal Waters. *Hydrobiologia*. 460:185-193.
- Mallin, M.A., K.E. Williams, E.C. Esham and R.P. Lowe. 2000. Effect of Human Development on Bacteriological Water Quality in Coastal Watersheds. *Ecological Applications*. 10(4):1047-1056.
- Weiskel, P., Howes, B. and G. Heufelder. 1996. Coliform Contamination of a Coastal Embayment: Sources and Transport Pathways. *Environmental Science and Technology*. 30(6):1872-1881.
- White, N.M., D.E. Line, J.D. Potts, W. Kirby-Smith, B. Doll and W.F. Hunt. 2000. Jump Run Creek Shellfish Restoration Project. *Journal of Shellfish Research*. 19(1):473-476.

Acknowledgments

Our appreciation to USEPA for financial assistance and to members of the project advisory committee, including Harriet Beale, Robin Downey, Bob Woolrich, Tim Determan, Marina Alberti, Marcie Bidwell, Mindy Roberts, Liz Lyman, Scott Clark, Kevin Dragon, Paul Williams, Chris May, Stuart Whitford, Shawn Ultican, Sylvia Goodwin, Jeff Chalfant and Chris Parsons.

For Further Information

Contact Stuart Glasoe of the Puget Sound Action Team at 360-725-5449 or sglasoe@psat.wa.gov. Full report available on-line at www.psat.wa.gov/Programs/shellfish/sf_lit_review0604.pdf.